

# Karyotype and sex chromosome differentiation in two *Nalassus* species (Coleoptera, Tenebrionidae)

Dirim Şendoğan<sup>1</sup>, Nurşen Alpagut-Keskin<sup>1</sup>

<sup>1</sup> Ege University, Faculty of Science, Department of Zoology, Section of Biology, Bornova, Izmir 35100 TURKEY

Corresponding author: Nurşen Alpagut-Keskin ([nursen.alpagut@ege.edu.tr](mailto:nursen.alpagut@ege.edu.tr))

Academic editor: D. Lachowska | Received 9 June 2016 | Accepted 2 August 2016 | Published 2 September 2016

<http://zoobank.org/CF663AE6-57EC-4C8D-BBF2-E8674B7B7475>

**Citation:** Şendoğan D, Alpagut-Keskin N (2016) Karyotype and sex chromosome differentiation in two *Nalassus* species (Coleoptera, Tenebrionidae). Comparative Cytogenetics 10(3): 371–385. doi: 10.3897/CompCytogen.v10i3.9504

## Abstract

Cytogenetic features of *Nalassus bozdagus* Nabozhenko & Keskin, 2010 and *Nalassus plebejus* Küster, 1850 were analysed using conventional and differential staining. Mitotic and meiotic chromosomal analysis revealed the diploid number as  $2n = 20 (9+X_{Yp})$  in both species. Besides the general resemblance of two *Nalassus* Mulsant, 1854 karyotypes, important differences related to variations in the number of meta-centric/submetacentric chromosomes, localization of highly impregnated regions which are considered as NOR and heterochromatin distribution are clearly observed. The most prominent difference between two species is found related to the X chromosome which is clearly larger in *N. bozdagus* and has a conspicuous secondary constriction on the long arm. As a result of silver staining, the existence of highly impregnated areas associated with  $X_{Yp}$  of *N. bozdagus* in both prophase I and metaphase I, suggests that NORs are seemingly located on sex chromosomes. On the other hand, the potential NORs of *N. plebejus* were observed only in prophase I nuclei. With the application of fluorescence dye DAPI, the AT rich chromosome regions and  $X_{Yp}$  which forms the parachute configuration were shown in both species.

## Keywords

Karyotype, Tenebrionidae, Helopini, *Nalassus bozdagus*, *Nalassus plebejus*, sex chromosomes, heterochromatin, NOR, DAPI



## Introduction

In the light of fossil and molecular dating analysis, the darkling beetles are dispersed and diversified over the last 180 million years prior to Gondwanan fragmentation. Tenebrionids represent a hyperdiverse family of Coleoptera with ca. 20000 recognized species worldwide. In consequence of undergoing multiple evolutionary radiations, tenebrionids show considerable morphological variations and several adaptations in life history traits such as feeding behaviour, habitat preferences, flight ability etc. Although higher level of tenebrionid phylogeny based on sequences from seven out of nine subfamilies shows well supported monophyly, the subfamilies Diaperinae, Pimeliinae and Tenebrioninae were recovered as paraphyletic or polyphyletic (Kergoat et al. 2014a, b).

The karyotypes of more than 250 darkling beetle species have been determined (Holecová et al. 2008a, Juan and Petitpierre 1991a, Blackmon and Jeffery 2015, Gregory 2016). Although most species present a karyotype with  $2n = 20$ , the diploid number ranges from  $2n = 14$  to  $2n = 38$  (Juan and Petitpierre 1991a, Pons 2004, Holecová et al. 2008a, Lira-Neto et al. 2012). Chromosomal data are only available for several representatives of subfamilies Lagriinae, Tenebrioninae, Pimelinae, Alleculinae and Diaperinae mostly distributed in Mediterranean (Juan and Petitpierre 1991a).

The genus *Nalassus* Mulsant, 1854 (Tenebrioninae: Helopini) comprises 71 described taxa distributed mainly in the Western Palearctic, but with disjunctively isolated species in the Russian Far East and Northern China (Medvedev 1987, Nabozhenko 2001). Even though a significant part of the species is found in alpine and subalpine mountainous belts with high level of local endemism, some species that are adapted to lowlands have wider distribution. In the recent reviews of *Nalassus* species from the European part of CIS (Commonwealth of Independent States), Caucasus, Iran, Georgia, China and Turkey, several new species and combinations were also noted (Nabozhenko 2001, Nabozhenko 2007, Nabozhenko 2008, Keskin and Nabozhenko 2010, Nabozhenko 2011, Nabozhenko 2013, Nabozhenko 2014, Nabozhenko and Ivanov 2015). Therefore, the actual diversity is certainly higher than previously estimated and the monophyly of the genus *Nalassus* needs to be tested with several new additional characters. The chromosomes of *Nalassus* have not yet been studied. Furthermore, cytogenetic data concerning the tribe Helopini which provide no more than chromosome numbers and sex determination systems are only known for some *Nesotes* and *Probatiscus* species (Juan and Petitpierre 1986, 1989, 1991a, 1991b, Palmer and Petitpierre 1997).

In this study, with the aim of providing first cytogenetic information about *Nalassus*, the mitotic and meiotic chromosomes of endemic *N. bozdagus* and widespread *N. plebejus* were analysed using conventional, DAPI fluorochrome staining and silver impregnation.

## Material and methods

The meiotic and mitotic chromosomes of 12 male *N. plebejus* and 4 male *N. bozdagus* individuals from Western Anatolia were analysed. The specimens of *Nalas-*



*sus plebejus* were retrieved from Ballıkayalar Natural Park, Gebze (40°50'22.96"N / 29°30'56.11"E, 120m) and the specimens of *Nalassus bozdagus* were collected from Bozdağ, İzmir (38°15'17.49"N / 27°57'44.72"E, 2300m). Adult beetles were collected on the trunks of trees and on the ground at night when they are active.

The chromosome preparations were obtained from the gonads of male specimens using Murakami and Imai's (1974) splashing method with some modifications. Briefly, testes were carefully dissected and macerated with sterilized needles. Testes were treated with hypotonic solution (0.65% KCl) for 5 minutes and fixed in 3:1 ethanol:acetic acid at least for 1 h on ice.

We also applied a microspreading method (Chandley 1994) for obtaining prophase I nuclei. The slides were stained with 4% Giemsa in phosphate buffer pH 6.8, for 20 minutes for standard staining. The silver impregnation technique of Patkin and Sorokin (1984) was performed to determine the possible NOR regions. Briefly, slides were incubated in distilled water for 30 min. at room temperature and stained with AgNO<sub>3</sub> working solution (2:1, 50% AgNO<sub>3</sub>: 2% gelatin containing 0.5% formic acid) in a humid chamber at 60 °C for 3-10 minutes. After a golden-brown color has developed, the reaction was stopped by rinsing with distilled water. Slides were then dehydrated, counterstained with 4% Giemsa in phosphate buffer pH 6.8.

For determining of heterochromatin distribution, the slides were mounted with antifade mounting medium with fluorochrome DAPI (Vectashield) specific to AT-rich chromosomal regions. The visualization of DAPI stained plates were carried out with Olympus BX50 fluorescent microscope.


The mitotic and meiotic plates were analysed and photographed with Zeiss Axio Scope light microscope using ZEN software. The chromosomal measurements were made with the LEVAN plugin (Sakamoto and Zacaro 2009) and the karyotypes and idiograms were created with the CHIAS plugin (Kato et al. 2011) of the programme IMAGE J (Rasband 1997-2015).

## Results

### Conventional Giemsa staining

Analysis of spermatogonial cells of *Nalassus bozdagus* and *Nalassus plebejus* revealed the diploid chromosome number as  $2n = 20 (9+X_{Y_p})$  (Fig. 1). In both species, most of the autosomes showed metacentric morphology, the X chromosomes were submetacentric and the y chromosomes were subtelocentric. In *N. bozdagus* the autosomal pairs 8 and 9 were submetacentric while in *N. plebejus* autosomal pairs 1, 5 and 8 were submetacentric. The smallest chromosome in both species was determined to be the y chromosome (~1 µm). The biggest chromosome of *Nalassus bozdagus* was the X chromosome (~4.315 µm), in *Nalassus plebejus* the biggest chromosome was the 1. chromosome with the length of 4.442 µm (Table 1).






**Figure 1.** a–b *N. bozdagus* 2n = 20. **a** karyotype **b** idiogram **c–d** *N. plebejus* 2n = 20 **c** karyotype **d** idiogram. Bar = 5 µm.

**Table 1.** Chromosome morphologies and measurements of *N. bozdagus* and *N. plebejus*. CI: centromere index, RL: relative length, AR: arm ratio, \*secondary constriction.

	<i>N. bozdagus</i>					<i>N. plebejus</i>				
Chromosome	Length (µ)	CI	%RL	AR	Morphology	Length (µ)	CI	%RL	AR	Morphology
1	3.895	46	12.4	1.20	m	4.442	35	14.26	1.8	sm
2	3.420	47	10.89	1.23	m	4.207	48	13.5	1.04	m
3	3.375	46	10.74	1.29	m	3.316	43	10.64	1.4	m
4	3.204	46	10.2	1.10	m	3.117	43	10	1.28	m
5	2.876	47	9.16	1.20	m	3.222	28	10.34	2.52	sm
6	2.715	48	8.64	1.06	m	3.040	44	9.76	1.27	m
7	2.204	48	7.02	1.07	m	2.439	45	7.83	1.18	m
8	2.162	33	6.88	2.13	sm	2.476	29	7.95	2.38	sm
9	2.149	32	6.84	2.01	sm	1.853	46	5.95	1.17	m
X	4.315	28	13.74	2.47	sm*	2.04	30	6.55	2.31	sm
y	1.097	20	3.5	3.88	st	1.010	18	3.24	1.17	st






**Figure 2.** **a–b** *N. bozdagus* with dark heterochromatic blocks on all chromosomes. **a** leptotene–zygotene **b** pachytene **c–d** *N. plebejus* with two distinctive heterochromatic blocks (black arrows); **c** leptotene–zygotene **d** pachytene. Orange arrow indicates  $X_{yp}$  sex bivalent, Bar = 5  $\mu\text{m}$ .

In prophase I nuclei, all chromosomes of *Nalassus bozdagus* showed dark heterochromatic blocks mainly located in centromeric regions (Fig. 2a–b). But in *Nalassus plebejus*, while most of the chromosomes have relatively small amounts of heterochromatin dispersed throughout the whole length (Fig. 2c–d), only 2 chromosomes with distinctive heterochromatic blocks were observed.


In diplotene/diakinesis of *N. plebejus*, 5–6 rod-shaped (terminal chiasma), 2–3 ring-shaped (two terminal chiasmata) and 1–2 cross-shaped (interstitial chiasma) bivalents were observed (Fig. 3a). In diakinesis/metaphase I; most of the homologous chromosomes of both species formed rod shaped bivalents due to being monochiasmatic and 2–3 ring shaped bivalents due to being bichiasmatic (Fig. 3b–c). In metaphase I plates, the parachute formation of sex bivalents was clearly observed for both *Nalassus* species (Fig. 4a–b). In metaphase II plates, relatively small sized 10 chromosomes (Fig. 4c–d) were observed. However, the plates which possessed the minute y chromosome were seemed to have only 9 chromosomes in their haploid sets. (Fig. 4d).

Sex chromosomes of two species were differed from each other by the length of X chromosome. The X chromosome of *N. bozdagus* was determined to be almost twice the size of the X chromosome of *N. plebejus* (Fig. 1, Table 1).





**Figure 3.** **a** diplotene–diakinesis in *N. plebejus* **b–c** diakinesis–metaphase I **b** *N. bozdagus* **c** *N. plebejus*. Bar = 5  $\mu$ m.



**Figure 4.** **a–b** metaphase I **a** *N. bozdagus* **b** *N. plebejus* **c–d** metaphase II **c** *N. bozdagus* **d** *N. plebejus*. Arrows show  $X_{yp}$  sex bivalents, Bar = 5  $\mu$ m.




## Differential staining

Silver nitrate staining of the chromosomes of *Nalassus bozdagus* revealed the presence of a highly impregnated nucleolus (NOR) associated with one of the long chromosomes in prophase I nuclei (Fig. 5a–b) and that  $X_{Y_p}$  sex bivalent is strongly argyrophilic in metaphase I (Fig. 5c). In *Nalassus plebejus*, these strongly argyrophilic regions were observed only in pachytene nuclei (Fig. 5d). With base-specific (A-T) DAPI staining; metaphase I plates and prophase I nuclei were observed. In metaphase I stages there were no significant difference between species (Fig. 6a–b). Prophase I nuclei of *N. bozdagus* showed strong signals on pericentromeric heterochromatic blocks compared to other chromosomal regions (Fig. 6c). On the other hand, in *N. plebejus* only some centromeric regions showed slightly stronger fluorescence signals (Fig. 6d).


## Discussion

The family Tenebrionidae is considered a karyologically conservative group due to the frequent occurrence of  $2n = 20$  formula (Juan and Petitpierre 1991a, Palmer and Petitpierre 1997). Heretofore, variation in the diploid chromosome numbers between



**Figure 5.** Silver nitrate staining **a–c** *N. bozdagus* **d** *N. plebejus*. Black arrows indicate NOR, red arrows indicate argyrophilic sex bivalents, Bar = 5  $\mu$ m.





**Figure 6.** DAPI staining **a** metaphase I of *N. plebejus* **b** metaphase I of *N. bozdagus* **c** prophase I nucleus of *N. bozdagus* **d** prophase I nucleus of *N. plebejus*. Arrows show  $X_{Y_p}$  sex bivalents and arrowheads indicate heterochromatic regions, Bars = 5  $\mu$ m.

14–38 within the family are also noted (Juan and Petitpierre 1991a, Pons 2004, Holecová et al. 2008a, Lira-Neto et al. 2012). Although, tenebrionid karyotypes characterized with predominant presence of metacentric chromosomes (Guenin 1950, 1951a, b; Smith 1952, Yadav and Pillai 1974, Yadav et al. 1980, Juan and Petitpierre 1988, 1989, 1990, Juan et al. 1989), several species from different subfamilies have mostly subtelocentric/acrocentric sets (e.g. *Laena reiteri* Weise 1877,  $2n = 18$ , Holecová et al. 2008a, *Palembus dermestoides* Fairmaire 1893,  $2n = 20$ , Almeida et al. 2000). Furthermore, many tenebrionid beetles possess similar chromosome number but differ in karyotype structure, which reveal additional evidence for karyotype divergence through the intra-chromosomal rearrangements. The major patterns of karyological variations in tenebrionid beetles are mainly observed in sex determining systems, chromosome



morphology and distribution of heterochromatin (Juan and Petitpierre 1990, Petitpierre et al. 1991, Juan and Petitpierre 1991a-b, Juan et al. 1993, Bruvo-Madaric et al. 2007).

Tenebrionidae comprises 9 subfamilies but most of the cytogenetically studied species (96%) belong to the Pimeliinae and Tenebrioninae (Bouchard et al. 2005, Holecová et al. 2008a). The diploid number in Pimeliinae shows a decrease from  $2n = 20$  to  $2n = 18$  caused by fusion of an autosomal pair while in Tenebrioninae there is a tendency of increased diploid number probably caused by centric fissions (Juan and Petitpierre 1991a).

We showed here that the karyotypes of *Nalassus bozdagus* and *Nalassus plebejus* consist of 10 pairs of chromosomes ( $2n = 20$ ) (Fig. 1), which is considered as modal chromosome number for Tenebrionidae (Juan and Petitpierre 1991a, Pons 2004, Holecová et al. 2008a, Lira-Neto et al. 2012). The presence of heteromorphic sex chromosomes for both species is confirmed by occurrence of a  $X_{Y_p}$  configuration in both conventionally (Fig. 4a-b) and differentially (Fig. 5c, 6a-b) stained metaphase I plates. The  $X_{Y_p}$  sex determining system is the most frequent type among Tenebrionidae as well as order Coleoptera (Smith and Virkki 1978, Juan and Petitpierre 1991a). However, sex chromosomes or sex determining systems are mentioned as one of the major chromosomal changes involved in tenebrionid divergence.

Besides the general resemblance of two *Nalassus* karyotypes, important differences related to X chromosomes, variations in the number of metacentric/submetacentric chromosomes (Fig. 1, Table 1), localization of highly impregnated regions which are considered as NOR (Fig. 5a-d) and heterochromatin distribution (Fig. 6c-d) are clearly observed. The most prominent difference between two species is found related to X chromosome which is clearly larger (13.74% of total complement) in *N. bozdagus* and has a conspicuous secondary constriction on the long arm (Fig. 1). It was also observed that metaphase I plates of *N. bozdagus* have relatively larger  $X_{Y_p}$  (Fig. 4a-b). The increase in relative length of X which does not alter parachute configuration is named as giant  $X_{Y_p}$  and generally thought to be derived from either heterochromatin amplification or translocation (Dutrillaux and Dutrillaux 2009). Difference in size and heterochromatin content of X chromosomes also observed in two closely related tenebrionid species of *Gonocephalum* Solier 1834 (Tenebrioninae) (Juan and Petitpierre 1989).

The differences found in chromosome morphology (1., 5. and 9. pairs) between these two *Nalassus* species are thought to be related to pericentromeric inversions that resulted in centromeric shift. Pericentromeric rearrangements are already known within several Coleopteran families such as Cicindelidae, Chrysomelidae, Meloidae, Scarabaeidae and Tenebrionidae (Serrano 1981, Petitpierre 1983, Juan et al 1990, Almeida et al 2000, Petitpierre and Garneria 2003, Wilson and Angus 2005, De Julio et al 2010, Petitpierre 2011).

The karyotypes of *N. bozdagus* and *N. plebejus* also show obvious differences, especially in distribution of heterochromatin. The presence of strong signals on pericentromeric heterochromatin blocks on all chromosomes of *N. bozdagus* (Fig. 2a-b) and only few chromosomes in *N. plebejus* (Fig. 2c-d) was supported with both conven-



tionally and differentially stained prophase I nuclei (Fig. 4g–h). Although, occurrence of heterochromatin observed mainly in the pericentromeric areas of the tenebrionid chromosomes, variability of heterochromatin localization and composition were also reported (Juan and Petitpierre 1989, 1991, Pons 2004, Rozek et al. 2004, Cabral-de-Mello et al. 2010, Schneider et al. 2007).

As a result of silver staining, the existence of highly impregnated areas associated with  $X_{Y_p}$  of *N. bozdagus* in both prophase I and metaphase I, suggests that NORs are seemingly located on sex chromosomes (Fig. 5a–c). On the other hand, the potential NORs of *N. plebejus* were observed only in prophase I nuclei (Fig. 5d). Similar findings on argyrophilic  $X_{Y_p}$  in metaphase I as well as prophase I were previously reported for several beetles such as *Zophobas* aff. *confusus* Gebien 1906 (Tenebrionidae) (Lira- Neto et al. 2012), *Lagria villosa* Fabricius 1781 (Tenebrionidae) (Goll et al. 2013), *Palembus dermestoides* (Tenebrionidae) and *Epicauta atomaria* Germar 1821 (Meloidae) (Almeida et al. 2000). Although, rDNA-FISH studies has shown that these strongly argyrophilic areas in prophase I bivalents are associated with NOR (Juan et al. 1993, Bruvo-Madaric et al. 2007), the existence of highly impregnated areas on sex chromosomes until metaphase I thought to be related to association and segregation of sex bivalents due to nucleolar material or distinctive heterochromatin (Juan et al. 1993, John and Lewis 1960). The association between sex chromosomes and nucleolar material is widely known for several animal groups from mammals to insects (Smith and Virkki 1978, Virkki et al. 1991, Tres 2005), although autosomal localization of NORs by FISH using 18S rDNA probes were also reported for some tenebrionid species (Goll et al. 2003, Juan et al. 1993).

It was observed that bichiasmatic autosomes form ring bivalent while monochiasmatics form rod bivalents due to terminal chiasmata (Fig. 3b–c). Ring bivalents are frequent among Tenebrionidae, Scarabaeidae, Meloidae, Buprestidae, Curculionidae, Chrysomelidae and Cerambycidae (Petitpierre 1985, Bisoi and Patnaik 1988, Petitpierre and Garneria 2003, Karagyan et al. 2004, 2012, Lachowska et al. 2004, 2006a, 2006b, Rozek et al. 2004, Angus et al. 2007, Holecová et al. 2008b). During diploten-diakinesis of *N. plebejus*, in addition to ring and rod bivalents we also observed cross shaped bivalents (Fig. 3a) due to interstitial chiasmata.

The information acquired from metaphase I plates of only few *Nesotes* Allard 1876 species (Juan and Petitpierre 1986, 1989, 1991a, 1991b) and *Probaticus ebeninus* A. Villa and J. B. Villa, 1838 (Palmer and Petitpierre 1997) are the only cytogenetic data concerning the tribe Helopini. On the basis of metaphase I plates, it was only briefly noted that 5 *Nesotes* species have similar  $2n = 20, 9 + X_{Y_p}$  formula (Juan and Petitpierre 1986, 1989, 1991a) and *P. ebeninus* have  $2n = 20, 9 + XY$  (Palmer and Petitpierre 1997). Although, our findings for chromosome numbers correspond to previous cytogenetic data, comparative genomic analyses of Helopini require detailed descriptions of chromosome morphologies.

In conclusion, this study revealed that the cytogenetic features differed between endemic *N. bozdagus* and widespread *N. plebejus*. But, in the absence of molecular cytogenetic and phylogenetic approaches, it is not possible to make a strong conclusion



about the major forces underlying these chromosomal variations. For definitive testing of the general trends in both *Nalassus* and tenebrionid karyotype evolution, it is necessary to increase the taxa sampling for major tenebrionid lineages.

## Acknowledgements

We are sincerely grateful to Dr. Maxim Nabozhenko and Dr Bekir Keskin for sharing their experience about Helopini. We also wish to thank to Dr Ersen Aydın Yağmur and Beril Gündoğan, for valuable help with collection of the specimens, Molecular Cytogenetic Lab members in Faculty of Medicine-Ege University for their help and Prof. Dr. R. Angus for his suggestions on earlier versions of this paper. Specimens included in this study are part of collected and sequenced material of a molecular systematic research project (TBAG-Project No:112T445).

## References

- Almeida MC, Zacaro AA, Cella DM (2000) Cytogenetic analysis of *Epicauta atomaria* (Meloidae) and *Palembus dermestoides* (Tenebrionidae) with Xy sex determination system using standard staining, C-bands, NOR and synaptonemal complex microspreading techniques. *Hereditas* 133: 147–157. doi: 10.1163/22119434-900000220
- Angus RB, Wilson CJ, Mann DJ (2007) A chromosomal analysis of 15 species of Gymno-pleurini, Scarabaeini and Coprini (Coleoptera: Scarabaeidae). *Tijdschrift Voor Entomologie* 150: 201–211.
- Bisoi MR, Patnaik SC (1988) A Chromosome Study of Seven Species of Indian Coleoptera (Meloidae, Tenebrionidae and Coccinellidae). *Caryologia* 41(3-4): 309–321. doi: 10.1080/00087114.1988.10797871
- Blackmon H, Jeffery PD (2015) Coleoptera Karyotype Database. *The Coleopterists Bulletin* 69.1: 174–175. doi: 10.1649/0010-065X-69.1.174
- Bouchard P, Lawrence JF, Davies AE, Newton AF (2005) Synoptic classification of the world Tenebrionidae (Insecta: Coleoptera) with a review of family-group names. *Annales Zoologici* 55(4): 499–530.
- Bruvo-Madaric B, Plohl M, Ugarkovic D (2007) Wide distribution of related satellite DNA families within the genus *Pimelia* (Tenebrionidae). *Genetica* 130: 35–42. doi: 10.1007/s10709-006-0017-2
- Cabral-de-Mello DC, Moura RC, Carvalho R, Souza MJ (2010) Cytogenetic analysis of two related *Deltochilum* (Coleoptera, Scarabaeidae) species: diploid number reduction, extensive heterochromatin addition and differentiation. *Micron* 41: 112–117. doi: 10.1016/j.micron.2009.10.005
- Chandley AC, Speed RM, Ma K (1994) Meiotic Chromosome Preparation. In: Gosden JR (Ed.) *Chromosome Analysis Protocols. Methods in Molecular Biology* 29: 27–40. doi: 10.1385/0-89603-289-2:27



- De Julio M, Rodrigues-Fernandes F, Costa C, Almeida MC, Cella DM (2010) Mechanisms of karyotype differentiation in Cassidinae sensu lato (Coleoptera, Polyphaga, Chrysomelidae) based on seven species of the Brazilian fauna and an overview of the cytogenetic data. *Micron* 41: 26–38. doi: 10.1016/j.micron.2009.07.013
- Dutrillaux AM, Dutrillaux B (2009) Sex chromosome rearrangements in Polyphaga beetles. *Sexual Development* 3(1): 43–54. doi: 10.1159/000200081
- Dutrillaux AM, Xie H, Dutrillaux B (2007) Nucleolus and chromosome relationships at pachynema in four Scarabaeoidea (Coleoptera) species with various combinations of NOR and sex chromosomes. *Chromosome Research* 15: 417–427. doi: 10.1007/s10577-007-1133-2
- Goll LG, Artoni RF, Vicari MR, Nogaroto V, Petitpierre E, Almeida MC (2013) Cytogenetic analysis of *Lagria villosa* (Coleoptera, Tenebrionidae): Emphasis on the mechanism of association of the Xy(p) Sex Chromosomes. *Cytogenetic and Genome Research* 139(1): 29–35. doi: 10.1159/000341674
- Gregory TR (2016) Animal Genome Size Database. <http://www.genomesize.com>
- Guenin HA (1950) Chromosomes et Hétérochromosomes de Ténébrionidés. *Genetica* 25: 157–182. doi: 10.1007/BF01784829
- Guenin HA (1951a) La formule chromosomiale de Coleopteres tenebrionides nordafricains, I. Pimeliines et Tentyriines. *Bulletin de la Société Vaudoise des Sciences Naturelles* 65: 7–18.
- Guenin HA (1951b) La formule chromosomiale de Coleopteres tenebrionides nordafricains, II. Erodiines. *Revue Suisse de Zoologie* 58: 471–475.
- Holecová M, Rozek M, Lachowska D (2008a) The First Cytogenetic Report on *Laena reitteri* Weise, 1877 with Notes on Karyotypes of Darkling Beetles. *Folia Biologica (Kraków)* 56(3-4): 213–217. doi: 10.3409/fb.56
- Holecová M, Rozek M, Lachowska D (2008b) Karyotype Analysis of Three Weevil Species from the Genera *Brachysomus* (Boheman, 1845) and *Strophosoma* Billberg, 1820 (Coleoptera, Curculionidae). *Folia Biologica (Kraków)* 56(1): 103–110. doi: 10.3409/fb56\_1-2.25-29
- John B, Lewis KR (1960) Nucleolar controlled segregation of the sex chromosomes in beetles. *Heredity* 15: 431–439. doi: 10.1038/hdy.1960.107
- Juan C, Petitpierre E (1986) Karyological Analyses on Tenebrionid beetles from Balearic Islands. *Génétique Ibérica* 38: 231–244.
- Juan C, Petitpierre E (1988) A chromosome survey of North African and Eastern Mediterranean tenebrionids (Coleoptera). *Cytobios* 54: 85–94.
- Juan C, Petitpierre E (1989) C-banding and DNA content in seven species of Tenebrionidae (Coleoptera). *Genome* 32: 834–839. doi: 10.1139/g89-519
- Juan C, Petitpierre E (1990) Karyological differences among Tenebrionidae (Coleoptera). *Genetica* 80: 101–108. doi: 10.1007/BF00127130
- Juan C, Petitpierre E (1991a) Chromosome numbers and sex determining systems in Tenebrionidae. *Advances in Coleopterology* 167–176.
- Juan C, Petitpierre E (1991b) Evolution of Genome Size in Darkling Beetles. *Genome* 34: 169–73. doi: 10.1139/g91-026
- Juan C, Petitpierre E, Oromi P (1989) Chromosomal Analyses on Tenebrionids from Canary Islands. *Cytobios* 57: 33–41.



- Juan C, Pons J, Petitpierre E (1993) Localization of tandemly repeated DNA sequences in beetle chromosomes by fluorescent *in situ* hybridization. *Chromosome Research* 1: 167–174. doi: 10.1007/BF00710770
- Karagyan G, Kuznetsova VG, Lachowska DL (2004) New Cytogenetic Data on Armenian Buprestids (Coleoptera, Buprestidae) with a Discussion of Karyotype Variation within the Family. *Folia Biologica (Kraków)* 52: 151–158. doi: 10.3409/1734916044527601
- Kato S, Ohmido N, Fukui K (2011) CHIAS 4 ver 1.02.
- Kergoat GJ, Bouchard P, Clamens AL, Abbate JL, Jourdan H, Jabbour-Zahab R, Genson G, Soldati L, Condamine FL (2014a) Cretaceous environmental changes led to high extinction rates in a hyperdiverse beetle family. *BMC Evolutionary Biology* 14(220): 1–13. doi: 10.1186/s12862-014-0220-1
- Kergoat GJ, Soldati L, Clamens AL, Jourdan H, Jabbour-Zahab R, Genson G, Bouchard P, Condamine FL (2014b) Higher level molecular phylogeny of darkling beetles (Coleoptera: Tenebrionidae). *Systematic Entomology* 39(3): 486–499. doi: 10.1111/syen.12065
- Keskin B, Nabozhenko MV (2010) A New Species and New Records of the Genus *Nalassus* Mulsant, 1854 (Coleoptera : Tenebrionidae : Helopini) from Turkey. *Annales Zoologici* 60(1): 23–28. doi: 10.3161/000345410X499489
- Lachowska D, Holecová M, Rozek M (2004) Notes on chromosome numbers and C-banding patterns in karyotypes of some weevils from Central Europe (Coleoptera, Curculionoidea: Apionidae, Nanophyidae, Curculionidae). *Folia Biologica* 52(1): 61–66.
- Lachowska D, Rozek M, Holecová M (2006a) Karyotypic characterization of three weevil species (Coleoptera: Curculionidae, Brachyderini). *Folia Biologica* 54(1): 13–17. doi: 10.3409/173491606777919139
- Lachowska D, Rozek M, Holecová M, Kajtoch L (2006b) Cytogenetic differences between *Peritelus familiaris* and *Centrinemus leucogrammus* (Coleoptera: Curculionidae: Entiminae: Peritelini). *European Journal of Entomology* 103: 687–690. doi: 10.14411/eje.2006.089
- Lira-Neto AC, Silva GM, Moura RC, Souza MJ (2012) Cytogenetics of the Darkling Beetles *Zophobas aff. confusus* and *Nyctobates gigas* (Coleoptera, Tenebrionidae). *Genetics and Molecular Research* 11(3): 2432–2440. doi: 10.4238/2012
- Medvedev GS (1987) Review of Tenebrionid Beetles of the Genus *Cylindronotus* Fald. (Coleoptera, Tenebrionidae) of Kazakhstan and Middle Asia. *Trudy Zoologicheskogo instituta AN SSSR* 170: 99–104. [In Russian]
- Murakami A, Imai H (1974) Cytological evidence for holocentric chromosomes of the silkworms, *Bombyx mori* and *B. mandarina*, (Bombycidae, Lepidoptera). *Chromosoma* 47: 167–178. doi: 10.1007/BF00331804
- Nabozhenko MV (2001) On the Classification of the Tenebrionid Tribe Helopini, with a Review of the Genera *Nalassus* Mulsant and *Odocnemis* Allard (Coleoptera, Tenebrionidae) of the European Part of CIS and the Caucasus. *Entomological Review* 81(8): 909–942.
- Nabozhenko MV (2007) Review of the subgenus *Helopondrus* Reitter, 1922 of the genus *Nalassus* Mulsant, 1854 (Coleoptera: Tenebrionidae) of Turkey. *Russian Entomological Journal* 16(4): 453–56.



- Nabozhenko MV (2008) *Nalassus glorificus* (Seidlitz, 1896) – New Synonym of *Nalassus pharnaces* Allard, 1876 (Coleoptera, Tenebrionidae). Caucasian Entomological Bulletin 4(1): 85.
- Nabozhenko MV (2011) Two New Species of the Genus *Nalassus* Mulsant, Subgenus *Helopondrus* Reitter (Coleoptera: Tenebrionidae) from Turkey. Stuttgarter Beitrage Zur Naturkunde A 4: 263–67.
- Nabozhenko MV (2013) Two New Taxa of the Genus *Nalassus* Mulsant, 1854 (Coleoptera: Tenebrionidae) from Georgia. Caucasian Entomological Bulletin 9(2): 261–64.
- Nabozhenko MV (2014) New Darkling Beetles of the Tribe Helopini (Coleoptera: Tenebrionidae) from Iran. Caucasian Entomological Bulletin 10(2): 237–41.
- Nabozhenko MV, Ivanov SN (2015) A New *Nalassus* Mulsant, 1854 (Coleoptera: Tenebrionidae: Helopini), the First Representative of the Genus from the Russian Far East. Zootaxa 3955(1): 137–41. doi: 10.11646/zootaxa.3955.1.9
- Palmer M, Petitpierre E (1997) New Chromosomal Findings on Tenebrionidae from Western Mediterranean. Caryologia 50(2): 117–23. doi: 10.1080/00087114.1997.10797391
- Patkin EL, Sorokin AV (1984) Nucleolus-Organizing Regions Chromosomes in Early Embryogenesis of Laboratory Mice. Bulletin of Experimental Biology and Medicine (USSR) 96: 92–94.
- Petitpierre E (1983) Karyometric Differences among Nine Species of the Genus *Chrysolina* Mots. (Coleoptera, Chrysomelidae). Canadian Journal of Genetics and Cytology 25: 33–39. doi: 10.1139/g83-006
- Petitpierre E (1985) Chromosome number and sex determining mechanism in four species of Galerucinae beetles with particular reference to the meiotic system of *Monolepta erythrocephala* (Ol.) (Chrysomelidae). Actas do II Congresso Ibérico de Entomologia (Lisboa) 1: 47–54.
- Petitpierre E (2011) Cytogenetics, cytotaxonomy and chromosomal evolution of Chrysomelinae revisited (Coleoptera, Chrysomelidae). ZooKeys 157: 67–79. doi: 10.3897/zookeys.157.1339
- Petitpierre E, Garnería I (2003) A cytogenetic study of the leaf beetle genus *Cyrtonus* (Coleoptera, Chrysomelidae). Genetica 119: 193–199. doi: 10.1023/A:1026010102779
- Petitpierre E, Juan C, Alvarez-Fuster A (1991) Evolution of Chromosomes and Genome Size in Chrysomelidae and Tenebrionidae. In: Belles X, Blas M, Zunino M (Eds) Advances in Coleopterology. European association of coleopterology, Barcelona, 129–44.
- Pons J (2004) Evolution of diploid chromosome number, sex-determining systems and heterochromatin in Western Mediterranean and Canarian species of the genus *Pimelia* (Coleoptera: Tenebrionidae). Journal of Zoological Systematics and Evolutionary Research 42: 81–85. doi: 10.1046/j.1439-0469.2003.00247.x
- Rasband WS (1997-2015) ImageJ, U.S. National Institutes of Health, Bethesda, Maryland, USA. <http://imagej.nih.gov/ij/>
- Rozek M, Lachowska D, Petitpierre E, Holecová M (2004) C-bands on chromosomes of 32 beetle species (Coleoptera: Elateridae, Cantharidae, Oedemeridae, Cerambycidae, Anthicidae, Chrysomelidae, Attelabidae and Curculionidae). Hereditas 140: 161–170. doi: 10.1111/j.1601-5223.2004.01810.x
- Sakamoto Y, Zacaro AA (2009) LEVAN, an ImageJ plugin for morphological cytogenetic analysis of mitotic and meiotic chromosomes, Initial version. <http://rsbweb.nih.gov/ij/>



- Schneider MC, Rosa SP, Almeida MC, Costa C, Cella DM (2007) Chromosomal similarities and differences among four Neotropical Elateridae (Conoderini and Pyrophorini) and other related species, with comments on the NOR patterns in Coleoptera. *Journal of Zoological Systematics and Evolutionary Research* 45: 308–316. doi: 10.1111/j.1439-0469.2006.00398.x
- Serrano J (1981) Chromosome numbers and karyotype evolution of Caraboidea. *Genetica* 55: 51–60. doi: 10.1007/BF00134005
- Smith SG (1952) The cytology of some tenebrionid beetles (Coleoptera). *Journal of Morphology* 91: 325–364. doi: 10.1002/jmor.1050910206
- Smith SG, Virkki N (1978) Coleoptera In: *Animal Cytogenetics Vol. 3 Insecta 5*. Borntraeger, Berlin.
- Tres L (2005) XY chromosomal bivalent: nucleolar attraction. *Molecular Reproduction and Development* 72: 1–6. doi: 10.1002/mrd.20334
- Virkki N, Mazzella C, Denton A (1991) Silver staining of the  $X_{Y_p}$  sex bivalent. *Cytobios* 67: 47–63.
- Wilson CJ, Angus RB (2005) A Chromosomal Analysis of 21 Species of Oniticellini and Onthophagini (Coleoptera: Scarabaeidae). *Tijdschrift Voor Entomologie* 148: 63–76. doi: 10.1163/22119434-900000167
- Yadav JS, Pillai RK (1974) Cytology of two species of Australian leafminers (Hispiinae Chrysomelidae). *Cytobios* 11: 75–79.
- Yadav JS, Pillai RK, Karamjeet (1980) Chromosome numbers of Tenebrionidae (Polyphaga: Coleoptera). *Biologia* 26: 31–41.